

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/21, 39/29, 39/39, C07K 14/16 // (A61K 39/29, 39:21)	A1	(11) International Publication Number: WO 99/51267 (43) International Publication Date: 14 October 1999 (14.10.99)
(21) International Application Number: PCT/CA99/00287 (22) International Filing Date: 1 April 1999 (01.04.99) (30) Priority Data: 09/055,744 7 April 1998 (07.04.98) US (71) Applicant (for all designated States except US): CONNAUGHT LABORATORIES LIMITED [CA/CA]; 1755 Steeles Avenue West, North York, Ontario M2R 3T4 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): SIA, Charles, D., Y. [CA/CA]; 27 Mabley Crescent, Thornhill, Ontario L4J 2Z7 (CA). CHONG, Pele [CA/CA]; 32 Estoril Street, Richmond Hill, Ontario L4C 0B6 (CA). KLEIN, Michel, H. [CA/CA]; 16 Munro Boulevard, Willowdale, Ontario M2P 1B9 (CA). (74) Agent: STEWART, Michael, I.; Sim & McBurney, 6th floor, 330 University Avenue, Toronto, Ontario M5G 1R7 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HIV-SPECIFIC CYTOTOXIC T-CELL RESPONSES		
(57) Abstract A method of generating an HIV-specific cytotoxic T-cell response in a host involves an initial administration of a T-helper molecule to the host to prime T-helper cells of the immune system of the host and a subsequent administration to the host of a mixture of the T-helper molecule and a T-cell inducing HIV-derived molecule to generate an HIV-specific T-cell response in the host.		